

$\pi\rho\omega$ VERTEX IN NUCLEAR MATTER

ANNA BARAN, AGNIESZKA BIENIEK, AND WOJCIECH BRONIEWSKI

*The H. Niewodniczański Institute of Nuclear Physics, ul. Radzikowskiego 152
PL-31342 Cracow, Poland*

Medium modifications of the $\pi\omega\rho$ vertex are analyzed in context of the $\omega \rightarrow \pi^0\gamma^*$ and $\rho \rightarrow \pi\gamma^*$ decays in nuclear matter. A relativistic hadronic model with mesons, nucleons, and $\Delta(1232)$ isobars is applied. A substantial increase of the widths for the decays $\omega \rightarrow \pi^0\gamma^*$ and $\rho \rightarrow \pi\gamma^*$ is found for photon virtualities in the range $0.3 - 0.6\text{GeV}$. This enhancement has a direct importance for the description of dilepton yields obtained in relativistic heavy-ion collisions.

[To appear in Proceedings of the 2nd International Symposium on Quantum Theory and Symmetries, Cracow, 18-21 July 2001, E. Kapuścik and A. Horzela, eds., World Scientific (Singapore, 2002)]

The experience gathered over the past years led to the conviction that the properties of hadrons in medium are substantially modified ^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20}. This view is supported by numerous studies of hadronic Green functions. Only a few investigations of the in-medium hadronic *couplings* exist ^{21,22,23,24,25,26,27}. For instance, it has been found that the value of the $\rho\pi\pi$ coupling is considerably enhanced. In Ref. ²⁸, which is the basis for this talk, we have examined the *in-medium* $\pi\omega\rho$ coupling. This vertex enters the decays $\omega \rightarrow \pi^0\gamma^*$ and $\rho \rightarrow \pi\gamma^*$, which are important for description of the dilepton production in relativistic heavy-ion collisions ^{29,30}. The existing calculations ^{31,32,33} have always used the unmodified (vacuum) value of the $\pi\omega\rho$ coupling. We have found that the medium effects can significantly alter the input of such calculations, thereby modifying the predicted dilepton production and creating hopes for a resolution of the long-standing problem of the experimental dilepton enhancement in the range $M_{l+l-} \sim 0.3 - 0.5\text{GeV}$.

In our study we use a typical hadronic model with meson, nucleon and Δ degrees of freedom, and for simplicity work at leading baryon density and at zero temperature. In addition, we constrain ourselves to the case where the decaying particle is at rest with respect to the medium. The effects of the relative motion turn out to be not very large ³⁴. In the vacuum the $\pi\omega\rho$ coupling is

$$-iV_{\pi\omega^\mu\rho^\nu} = i\frac{g_{\pi\omega\rho}}{F_\pi}\epsilon^{\mu\nu\rho Q}, \quad (1)$$




Figure 1. Diagrams included in our calculation (crossed ones not shown). Wavy lines denotes the ρ or ω , dashed lines the pions, solid lines the in-medium nucleon, and double lines the Δ resonance.

where we have used the notation $\epsilon^{\mu\nu p Q} = \epsilon^{\mu\nu\alpha\beta} p_\alpha Q_\beta$, and $F_\pi = 93\text{MeV}$ is the pion decay constant. In our convention Q is the incoming momentum of the pion, p is the outgoing momentum of the ρ , and $q = Q - p$ is the outgoing momentum of the ω . The vector mesons are, in general, virtual. For the kinematics where all external momenta vanish, the vector meson dominance models gives $g_{\pi \rightarrow \omega \rho} = -\frac{g_\rho g_\omega}{e^2} g_{\pi \gamma \gamma}$, with $g_{\pi \gamma \gamma} = \frac{e^2}{4\pi^2 F_\pi}$. The nucleon propagator in nuclear matter is decomposed into the *free* and *density* parts in the usual way ⁴:

$$iS(k) = iS_F(k) + iS_D(k) = i(\gamma^\mu k_\mu + m_N) \left[\frac{1}{k^2 - m_N^2 + i\varepsilon} + \frac{i\pi}{E_k} \delta(k_0 - E_k) \theta(k_F - |k|) \right]. \quad (2)$$

Here m_N denotes the nucleon mass, $E_k = \sqrt{m_N^2 + k^2}$, and k_F is the Fermi momentum. The Rarita-Schwinger propagator ^{35,36} for the spin 3/2 $\Delta(1232)$ particle has the form

$$iS_\Delta^{\mu\nu}(k) = i \frac{\gamma^\mu k_\mu + M_\Delta}{k^2 - M_\Delta^2} \left(-g^{\mu\nu} + \frac{1}{3} \gamma^\mu \gamma^\nu + \frac{2k^\mu k^\nu}{3M_\Delta^2} + \frac{\gamma^\mu k^\nu - \gamma^\nu k^\mu}{3M_\Delta} \right). \quad (3)$$

The effects of the non-zero width of the Δ are incorporated by replacing M_Δ by the complex mass $M_\Delta - i\Gamma/2$.

The diagrams used in our study are shown in Fig. 1. The density effects are produced when exactly one of the nucleon lines in each of the our diagrams of Fig. 1 involves the nucleon *density* propagator, S_D . The diagrams with

more than one S_D vanish for kinematic reasons. On the other hand, diagrams with no S_D describe the vacuum polarization effects, not considered in the present study devoted to the Fermi sea effect only.

The meson-baryon vertices needed in our analysis have the standard form

$$\begin{aligned} -iV_{\omega^\mu NN} &= ig_\omega \gamma^\mu, \\ -iV_{\rho_b^\mu NN} &= i\frac{g_\rho}{2}(\gamma^\mu - \frac{i\kappa_\rho}{2m_N}\sigma^{\mu\nu}p_\nu)\tau^b, \\ -iV_{\pi^a NN} &= \frac{g_A}{2F_\pi}\not{Q}\gamma^5\tau^a. \end{aligned} \quad (4)$$

For the interactions involving the Δ resonance we use

$$\begin{aligned} -iV_{N\Delta^\alpha\pi^a} &= g_{N\Delta\pi}Q^\alpha T^a, \\ -iV_{N\Delta^\alpha\rho_b^\mu} &= ig_{N\Delta\rho}(\not{p}\gamma^5 g^{\alpha\mu} - \gamma^\mu\gamma^5 p^\alpha)T^b, \\ -iV_{\Delta^\alpha\Delta^\beta\omega^\mu} &= -ig_\omega(\gamma^\alpha\gamma^\mu\gamma^\beta - \gamma^\beta g^{\alpha\mu} - \gamma^\alpha g^{\beta\mu} + \gamma^\mu g^{\alpha\beta}). \end{aligned} \quad (5)$$

Above T^a is the standard isospin $\frac{1}{2} \rightarrow \frac{3}{2}$ transition matrix^{37,36,38,39,40}. We choose the following values for the physical parameters: $g_\omega = 10.4$, $g_\rho = 5.2$, $g_A = 1.26$, $g_{N\Delta\pi} = 2.12/m_\pi$, $g_{N\Delta\rho} = 2.12\sqrt{2}/m_\pi$, $\kappa_\rho = 3.7$.

The $\pi\omega\rho$ vertex has the following tensor structure in the nuclear medium moving with the four-velocity u :

$$\begin{aligned} A^{\mu\nu} &= A_1\varepsilon^{\mu\nu pQ} + A_2\varepsilon^{\mu\nu uQ} + A_3\varepsilon^{\mu\nu pu} + A_4\varepsilon^{\mu\nu pQ}p^\nu \\ &+ A_5\varepsilon^{\mu\nu pQ}Q^\nu + A_6\varepsilon^{\mu\nu pQ}u^\nu + A_7\varepsilon^{\nu upQ}p^\mu + A_8\varepsilon^{\nu upQ}Q^\mu + A_9\varepsilon^{\nu upQ}u^\mu. \end{aligned} \quad (6)$$

This structure, restricted by the Lorentz invariance and parity, is more involved than in the vacuum case (1) due to the availability of the additional four-vector u . The leading-density calculation of diagrams of Fig. 1 is very simple, and follows the steps of Ref.²⁷. First, the trace factors are evaluated, then k^μ is substituted by $m_N u^\mu$, with $u^\mu = (1, 0, 0, 0)$ in the rest frame of the medium, and, finally, the loop integration is replaced by $\int \frac{d^4k}{(2\pi)^4}\Theta(k_f - |\vec{k}|)\delta(k_0 - E_k) \rightarrow \frac{1}{8\pi}\rho_B$, where ρ_B denotes the baryon density²⁷.

For the pion is at rest with respect to the medium, the four-vectors Q and u are parallel, and then $A^{\mu\nu} = (A_1 + A_3/m_\pi)\varepsilon^{\mu\nu pQ}$. We introduce the notation $A^{\mu\nu} = i\frac{e^2}{g_\rho g_\omega}\left(\frac{g_{\pi\rho\omega}}{F_\pi} + \frac{\rho_B}{F_\pi}B\right)\varepsilon^{\nu\mu pQ}$, which puts the medium effects in B . Our results below will be presented relative to the vacuum value $g_{\text{vac}} = g_{\pi\rho\omega}$, hence we define $g_{\text{eff}} \equiv g_{\pi\rho\omega} + \rho_0 B$.

For the processes $\omega \rightarrow \pi^0\gamma^*$ and $\rho^a \rightarrow \pi^a\gamma^*$, where ω and ρ are on mass shell, we use the following vacuum values of the coupling constants: $g_{\omega \rightarrow \rho\pi} =$




Figure 2. Top: $|g_{\text{eff}}/g_{\text{vac}}|^2$ for the decay $\omega \rightarrow \gamma^* \pi^0$ at the nuclear saturation density, plotted as a function of the virtuality of the photon, M . Bottom: the same for the decay ρ .

-1.13 and $g_{\rho \rightarrow \omega \pi} = -0.76$, obtained from the experimental partial decay width $\Gamma_{\omega \rightarrow \pi^0 \gamma} = 717 \text{keV}$ and $\Gamma_{\rho^a \rightarrow \pi^a \gamma} = 79 \text{keV}$. These coupling constants differ from each other, as well as from $g_{\pi \rightarrow \omega \rho} = -\frac{g_\rho g_\omega}{e^2} g_{\pi \gamma \gamma} = -1.36$ inferred from the π^0 decay. This difference is a manifestation of form-factor effects related to virtuality of vector mesons.

The numerical results are shown in Fig. 2. At low values of M the effective coupling constant remains practically unchanged. At higher values of M , above 0.2GeV , the value of g_{eff} is much larger than in the vacuum: for the square of g_{eff} we have an enhancement by a factor of ~ 2 for the ω , and ~ 5 for the ρ around $M = 0.4 \text{GeV}$. These values strongly indicate the possibility of large medium effects in the Dalitz decays of vector mesons. The enhancement of Fig. 2 directly affects the calculations of the dilepton yields in relativistic heavy-ion collisions. The theoretical output from the Dalitz decays

of vector mesons becomes enhanced in the region of $0.3 - 0.6\text{GeV}$, which is exactly where the existing calculations have serious problems in supplying enough strength to explain the CERES and HELIOS experiments^{29,30}. The dilepton production rate from the Dalitz decays of a vector meson is given by the formula³²

$$\frac{dN_{l+l-}}{d^4x dM^2} = \int \frac{d^3p_v}{(2\pi)^3 E_v} f(p_v) \frac{3m_v}{\pi M^3} \Gamma_{v \rightarrow \pi^0 \gamma^*} \Gamma_{\gamma^* \rightarrow l+l-} \quad (7)$$

where v relates to the vector meson, and $f(p_v)$ is the distribution function. The width $\Gamma_{v \rightarrow \pi^0 \gamma^*}$ depends on the baryon density, ρ_B , which in turn, depends on the space-time point x . Clearly, an increased value of $\Gamma_{\omega \rightarrow \pi^0 \gamma^*}$ results in an increased dilepton yield. A more exact estimate requires a detailed model of the evolution of the fireball, and is left for a further study³⁴. According to our analysis (*cf.* Fig. 3), at $M \approx 400\text{MeV}$ we have $\Gamma_{\omega \rightarrow \pi^0 \gamma^*} \approx 900\text{keV}$ and $\Gamma_{\rho^0 \rightarrow \pi^0 \gamma^*} \approx 250\text{keV}$. However, in a thermalized system due to isospin degeneracy the ρ mesons are three times more numerous, hence the effect of the ρ is comparable in size to the effect of the ω . The Dalitz yields from the π^0 decays are not altered by the medium for a double reason: first, virtually all pions decay outside of the fireball due to their long lifetime, second, the width for this decay is practically unaltered by the medium²⁸.

Acknowledgements

Research supported by the Polish State Committee for Scientific Research grant 2P03B09419.

References

1. G. E. Brown and M. Rho, *Phys. Rev. Lett.*, 66:2720, 1991.
2. L. S. Celenza, A. Pantziris, C. M. Shakin, and W.-D. Sun, *Phys. Rev.*, C45:2015, 1992.
3. B. D. Serot and J. D. Walecka, *Adv. Nucl. Phys.*, 16:1, 1986.
4. S. A. Chin, *Ann. Phys. (NY)*, 108:301, 1977.
5. T. Hatsuda, H. Shiomi, and H. Kuwabara, *Prog. Theor. Phys.*, 95:1009, 1996.
6. H.-C. Jean, J. Piekarewicz, and A. G. Williams, *Phys. Rev.*, C49:1981, 1994.
7. M. Herrmann, B. L. Friman, and W. Noerenberg, *Nucl. Phys.*, A560:411, 1993.
8. B. Friman and H. J. Pirner, *Nucl. Phys.*, A617:496, 1997.

9. R. Rapp, G. Chanfray, and J. Wambach, *Nucl. Phys.*, A617:472, 1997.
10. S. Leupold and U. Mosel, *Prog. Part. Nucl. Phys.*, 42:221, 1999.
11. S. Gao, C. Gale, C. Ernst, H. Stocker, and W. Greiner, *Nucl. Phys.*, A661:518, 1999.
12. T. Hatsuda and S. H. Lee, *Phys. Rev.*, C46:R34, 1993.
13. S.-H. Lee, *Nucl. Phys.*, A670:119, 2000.
14. U. Vogl and W. Weise, *Prog. Part. Nucl. Phys.*, 27:195, 1991.
15. F. Klingl, N. Kaiser, and W. Weise, *Nucl. Phys.*, A624:527, 1997.
16. G. Chanfray, *Nucl. Phys.*, A685:328, 2001.
17. W. Broniowski and B. Hiller, *Phys. Lett.*, B392:267, 1997.
18. V. L. Eletsky, B. L. Ioffe, and J. I. Kapusta, *Eur. J. Phys.*, A3:381, 1998.
19. B. Friman, *Acta Phys. Polon.*, B29:3195, 1998.
20. M. Lutz, B. Friman, and G. Wolf, *Nucl. Phys.*, A661:526, 1999.
21. C. Song and V. Koch, *Phys. Rev.*, C54:3218, 1996.
22. W. Broniowski, W. Florkowski, and B. Hiller, *Acta Phys. Polon.*, B30:1079, 1999.
23. W. Broniowski, W. Florkowski, and B. Hiller, *Eur. Phys. J.*, A7:287, 2000.
24. M. Urban, M. Buballa, R. Rapp, and J. Wambach, *Nucl. Phys.*, A641:433, 1998.
25. M. Urban, M. Buballa, R. Rapp, and J. Wambach, *Nucl. Phys.*, A673:357, 2000.
26. M. K. Banerjee and J. A. Tjon, *Phys. Rev.*, C56:497, 1997.
27. W. Broniowski, W. Florkowski, and B. Hiller, nucl-th/0103027.
28. A. Bielek, A. Baran, and W. Broniowski, nucl-th/0106053.
29. CERES Collab., G. Agakichiev *et al.*, *Phys. Rev. Lett.*, 75:1272, 1995.
30. HELIOS/3 Collab., M. Masera *et al.*, *Nucl. Phys.*, A590:93c, 1995.
31. G. Q. Li, C. M. Ko, and G. E. Brown, *Nucl. Phys.*, A606:568, 1996.
32. P. Koch, *Z. Phys.*, C57:283, 1993.
33. R. Rapp and J. Wambach, *Adv. Nucl. Phys.*, 25:1, 2000.
34. A. Bielek and W. Broniowski, to be published.
35. W. Rarita and J. Schwinger, *Phys. Rev.*, 60:61, 1941.
36. M. Benmerrouche, R. M. Davidson, and N. C. Mukhopadhyay, *Phys. Rev.*, C39:2339, 1989.
37. T. O. E. Ericson and W. Weise, *Pions and nuclei*, Clarendon Press, Oxford, 1988.
38. T. R. Hemmert, B. R. Holstein, and J. Kambor, *J. Phys. G*, G24:1831, 1998.
39. V. Pascalutsa and R. Timmermans, *Phys. Rev.*, C60:042201, 1999.
40. H. Haberzettl, nucl-th/9812043.